arXiv:1701.04589v1 [math.CA] 17 Jan 2017

ON FRACTIONAL KINETIC EQUATIONS AND THEIR SUMUDU TRANSFORM MULTIPARAMETER STRUVE FUNCTIONS BASED SOLUTIONS

K.S. NISAR*, F.B.M. BELGACEM, M. S. ABOUZAID

Abstract. This research paper treats fractional kinetic equations using the Sumudu transform operator. The exact solutions obtained are presented in terms of Struve functions of four parameters. By way of obtaining solutions some novel and useful and novel kinetic theorems are presented in light of the Sumudu properties. Results obtained in this study may be pragmatically used in many branches theoretical and experimental science applications, not the least of which are mathematical physics, and various engineering fields.

1. Introduction

The Struve function $H_v(z)$ and modified Struve function $L_v(z)$ are are given by the following infinite series, respectively,

$$H_v(z) = \left(\frac{z}{2}\right)^{v+1} \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+v+\frac{3}{2}\right)} \left(\frac{z}{2}\right)^{2k}, \tag{1.1}$$

and

$$L_v(z) = \left(\frac{z}{2}\right)^{v+1} \sum_{k=0}^{\infty} \frac{1}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+v+\frac{3}{2}\right)} \left(\frac{z}{2}\right)^{2k}. \tag{1.2}$$

The Struve function of order v (see [40]) is the solution of the non-homogeneous Bessel differential equation:

$$x^2y''(x) + xy'(x) + \left(x^2 - v^2\right) y(x) = \frac{4(x/2)^{v+1}}{\sqrt{\pi}\Gamma(v+1/2)}, \quad v \in \mathbb{C}. \tag{1.3}$$

where $\Gamma(z)$ is the gamma function and, $L_v(z)$ is related to $H_v(z)$, by the relation

$$L_n(z) = -ie^{-n\pi i/2} H_n(iz). \tag{1.4}$$

While, the homogeneous solutions of (1.3) are the Bessel functions, the particular solutions may be given to correspond to Struve functions. For complex parameter, v, modified

2010 *Mathematics Subject Classification.* 26A33, 44A10, 44A20, 33E12.

Key words and phrases. Fractional kinetic equations, Sumudu transforms, generalized Struve function, Fractional calculus .

*corresponding author.

Struve functions, L_v, turn out to be solutions of a modified version of equation (1.3), where the LHS zeroth coefficient is replaced by: $-(x^2+v^2)$. Applications of Struve functions can be found in various branches of applied science (see [1, 21, 29, 35, 36]).

Generalized versions of Struve function are done by extending its domain or expanding the type and number of parameters. In particular, a four- parameter generalized struve function, studied by Singh [37], is defined by:

$$H_{p,\mu}^{\lambda,\alpha}(x) := \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(\alpha k+\mu)\,\Gamma\left(\lambda k+p+\frac{3}{2}\right)} \left(\frac{x}{2}\right)^{2k+p+1}, p \in \mathbb{C} \tag{1.5}$$

where $\lambda > 0, \alpha > 0$ and μ an arbitrary parameter.

For subsequent need towards our objective below we also recall the generalized Mittag-Leffler function [24] defined by,

$$E_{\alpha,\beta}(x) = \sum_{n=0}^{\infty} \frac{x^n}{\Gamma(\alpha n+\beta)}. \tag{1.6}$$

The main motive of this paper is to study the solution of generalized form of the fractional kinetic equation involving generalized Struve function of four parameters with the help of Sumudu transform.

The Sumudu transform introduced by Watugala (see [41, 42]). For more details about Sumudu transform, see ([2–10, 13, 22]). The Sumudu transform over the set functions

$$A = \left\{ f(t) \,\middle|\, \exists\ M, \tau_1, \tau_2 > 0, |f(t)| < Me^{|t|/\tau_j}\ ,\ \text{if } t \in (-1)^j \times [0,\infty) \right\}$$

is defined by

$$G(u) = S[f(t); u] = \int_0^{\infty} f(ut)\, e^{-t} dt,\ u \in (-\tau_1, \tau_2). \tag{1.7}$$

The significance of fractional differential equations in the field of applied science increased more attention not only in mathematics but also in physics, dynamical systems, control systems and systems engineering, to create the mathematical model of numerous physical phenomena ([14–16, 18, 19, 25–28, 30–34, 43]. To carry our investigation,we would like to recall the following results due to Haubold and Mathai [20]. The fractional differential equation between rate of change of reaction was established by Haubold and Mathai [20]. The destruction rate and the production rate are given as follows

$$\frac{dN}{dt} = -d(N_t) + p(N_t) \tag{1.8}$$

where $N = N(t)$ the rate of reaction ,$d = d(N)$ the rate of destruction, $p = p(N)$ the rate of production and N_t denote the function defined by $N_t(t^*) = N(t - t^*), t^* > 0$

The special case of (1.8), for spatial fluctuations or in homogeneities in $N(t)$ the quantity are neglected, that is the equation

$$\frac{dN}{dt} = -c_i N_i(t) \tag{1.9}$$

with the initial condition that $N_i(t=0) = N_0$ is the number of density of species i at time $t = 0$ and $c_i > 0$. If we reject the index i and integrate the standard kinetic equation (1.9), we have

$$N(t) - N_0 = -c_0 D_t^{-1} N(t) \tag{1.10}$$

where ${}_0D_t^{-1}$ is the special case of the Riemann-Liouville integral operator ${}_0D_t^{-\upsilon}$ defined as

$${}_0D_t^{-\upsilon} f(t) = \frac{1}{\Gamma(\upsilon)} \int_0^t (t-s)^{\upsilon-1} f(s)\, ds, t > 0, \Re(\upsilon) > 0 \tag{1.11}$$

The fractional generalization of the standard kinetic equation (1.10) given by Haubold and Mathai [20] as:

$$N(t) - N_0 = -c_0^{\upsilon} D_t^{-1} N(t) \tag{1.12}$$

and obtained the solution of (1.12) as follows

$$N(t) = N_0 \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(\upsilon k + 1)} (ct)^{\upsilon k} \tag{1.13}$$

Further, Saxena and Kalla [34] considered the following fractional kinetic equation:

$$N(t) - N_0 f(t) = -c^{\upsilon} \left({}_0D_t^{-1} N\right)(t) \qquad (\Re(\upsilon) > 0) \tag{1.14}$$

where $N(t)$ denotes the number density of a given species at time t, $N_0 = N(0)$ is the number density of that species at time $t = 0$, c is a constant and $f \in L(0, \infty)$. By applying the Laplace transform to (1.14),

$$\begin{aligned} L[N(t)](p) &= N_0 \frac{F(p)}{1 + c^{\upsilon} p^{-\upsilon}} \\ &= N_0 \left(\sum_{n=0}^{\infty} (-c^{\upsilon})^n p^{-n\upsilon} \right) F(p) \qquad \left(n \in N_0, \left| \frac{c}{p} \right| < 1 \right), \end{aligned} \tag{1.15}$$

where the Laplace transform ([38]) is defined by

$$F(p) = L[f(t)] = \int_0^{\infty} e^{-pt} f(t)\, dt \quad \Re(p) > 0 \tag{1.16}$$

The Sumudu transform of (1.11) is (see, ([3]), p.106,eqn (2.1)) :

$$S\left[{}_0D_t^{-\upsilon} f(t)\right] = u^{\upsilon} G(u) \tag{1.17}$$

where $G(u)$ is defined in (1.7). It is clear that the Sumudu transform of $f(t) = t^{\alpha}$ is given by

$$S[f(t)] = \int_0^{\infty} (tu)^{\alpha} e^{-t} dt = u^{\alpha}\Gamma(\alpha+1), \Re(\alpha) > -1 \tag{1.18}$$

2. Solution of generalized fractional Kinetic equations

We devote this section to the derivative of the exact solution of time fractional kinetic equation via the well-known Sumudu transform. The results are as follows:

Theorem 1. *If $d > 0, \upsilon > 0, \alpha, c, l, t \in \mathbb{C}$,$\Re(l) > -1$ and $\Re(u) > 0$ with $|u| < d^{-1}$, then the solution of the equation*

$$N(t) - N_0 H_{l,\mu}^{\lambda,\alpha}(t) := -d^{\upsilon}\ {}_0D_t^{-\upsilon} N(t), \tag{2.1}$$

is given by the following formula

$$N(t) = \frac{N_0}{2} \sum_{k-0}^{\infty} \frac{(-1)^k \Gamma(2k+l+2)}{\Gamma(\alpha k+\mu)\Gamma\left(\lambda k+l+\frac{3}{2}\right)} \left(\frac{t}{2}\right)^{2k+l} E_{\upsilon,2k+l+1}(-d^{\upsilon}t^{\upsilon}) \tag{2.2}$$

where $E_{\upsilon,2k+l+1}(.)$ is the generalized Mittag-Leffler function given in (1.6).

Proof. Taking the Sumudu transform to the both sides of (2.1), we obtain the following relation,

$$S\{N(t)\} - N_0 S\left\{H_{l,\mu}^{\lambda,\alpha}(t); p\right\} = -d^{\upsilon} S\left\{{}_0D_t^{-\upsilon} N(t)\right\}$$

Now, using the definition given in (1.5), we get

$$\overset{*}{N}(u) - N_0 \left\{ S\left(\sum_{k-0}^{\infty} \frac{(-1)^k}{\Gamma(\alpha k+\mu)\Gamma\left(\lambda k+l+\frac{3}{2}\right)} \left(\frac{t}{2}\right)^{2k+l+1} \right) \right\} = -d^{\upsilon}u^{\upsilon}\overset{*}{N}(u)$$

where $\overset{*}{N}(u) = S\{N(t), u\}$ and $S\left\{t^{\lambda-1}\right\} = u^{\lambda-1}\Gamma(\lambda)$ gives,

$$\overset{*}{N}(u) - N_0 \sum_{k-0}^{\infty} \frac{(-1)^k}{\Gamma(\alpha k+\mu)\Gamma\left(\lambda k+l+\frac{3}{2}\right)} u^{2k+l+1}\Gamma(2k+l+2) = -d^{\upsilon}u^{\upsilon}\overset{*}{N}(u),$$

$$\overset{*}{N}(u)(1+d^{\upsilon}u^{\upsilon}) = N_0 \sum_{k-0}^{\infty} \frac{(-1)^k \left(\frac{u}{2}\right)^{2k+l+1} \Gamma(2k+l+2)}{\Gamma(\alpha k+\mu)\Gamma\left(\lambda k+l+\frac{3}{2}\right)}$$

$$\overset{*}{N}(u) = \frac{N_0}{(1+d^{\upsilon}u^{\upsilon})} \sum_{k-0}^{\infty} \frac{(-1)^k \left(\frac{u}{2}\right)^{2k+l+1} \Gamma(2k+l+2)}{\Gamma(\alpha k+\mu)\Gamma\left(\lambda k+l+\frac{3}{2}\right)}$$

$$= N_0 \sum_{k-0}^{\infty} \frac{(-1)^k \left(\frac{u}{2}\right)^{2k+l+1} \Gamma(2k+l+2)}{\Gamma(\alpha k+\mu)\Gamma\left(\lambda k+l+\frac{3}{2}\right)} \sum_{r-0}^{\infty} (-1)^r (du)^{\upsilon r}$$

Now, taking the inverse transform of the above expression and using the relation

$$S^{-1}\left\{u^{v}\right\}=\frac{t^{\lambda-1}}{\Gamma\left(v\right)},\quad \Re\left(u\right)>0,\Re\left(v\right)>0,$$

we get

$$\begin{aligned} N\left(t\right) &= N_{0}\sum_{k-0}^{\infty}\frac{\left(-1\right)^{k}\left(2\right)^{-(2k+l+1)}\Gamma\left(2k+l+2\right)}{\Gamma\left(\alpha k+\mu\right)\Gamma\left(\lambda k+l+\frac{3}{2}\right)}\sum_{r-0}^{\infty}\frac{\left(-1\right)^{r}d^{vr}t^{2k+vr+l}}{\Gamma\left(2k+l+1+vr\right)} \\ &= N_{0}\sum_{k-0}^{\infty}\frac{\left(-1\right)^{k}\Gamma\left(2k+l+2\right)t^{2k+l}}{\Gamma\left(\alpha k+\mu\right)\Gamma\left(\lambda k+l+\frac{3}{2}\right)\left(2\right)^{2k+l+1}}\sum_{r-0}^{\infty}\frac{\left(-1\right)^{r}d^{vr}t^{vr}}{\Gamma\left(2k+l+1+vr\right)} \end{aligned}$$

In view of (1.6), we obtain the desired result. □

Corollary 2.1. *If we put* $\alpha=\lambda=1$ *and* $\mu=3/2$ *in* (2.1) *then we obtain the solution of fractional kinetic equation involving Struve function* $H_{v}\left(z\right)$ *as :*

For $d>0,\upsilon>0,c,l,t\in\mathbb{C},\Re\left(l\right)>-1$ and $\Re\left(u\right)>0$, then the solution of the equation

$$N\left(t\right)-N_{0}H_{l,3/2}^{1,1}\left(t\right):=-d^{\upsilon}\ {}_{0}D_{t}^{-\upsilon}N\left(t\right),$$

is given by the following formula

$$N\left(t\right)=\frac{N_{0}}{2}\sum_{k-0}^{\infty}\frac{\left(-1\right)^{k}\Gamma\left(2k+l+2\right)}{\Gamma\left(k+l+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)}\left(\frac{t}{2}\right)^{2k+l}E_{v,2k+l+1}\left(-d^{\upsilon}t^{\upsilon}\right)$$

Theorem 2. *If* $d>0,\upsilon>0,\alpha,c,b,l,t\in\mathbb{C}$, $\Re\left(l\right)>-1$ *and* $\Re\left(u\right)>0$ *then for the solution of the equation*

$$N\left(t\right)-N_{0}H_{p,\mu}^{\lambda,\alpha}\left(d^{\upsilon}t^{\upsilon}\right)=-d^{\upsilon}{}_{0}D_{t}^{-\upsilon}N\left(t\right) \tag{2.3}$$

there holds the formula

$$\begin{aligned} N\left(t\right) &= N_{0}\left(\frac{d^{v}}{2}\right)^{l+1}t^{lv+v-1} \\ &\times\sum_{k=0}^{\infty}\frac{\left(-1\right)^{k}\Gamma\left(2kv+vl+\nu+1\right)}{\Gamma\left(\alpha k+\mu\right)\Gamma\left(\lambda k+l+\frac{3}{2}\right)}\left(\frac{d^{v}t^{v}}{2}\right)^{2k}E_{v,(2k+l+1)v}\left(-d^{\upsilon}t^{\upsilon}\right), \end{aligned} \tag{2.4}$$

where $E_{v,2kv+lv+v+1}\left(.\right)$ *is the generalized Mittag-Leffler function* (1.6).

Proof. The proof can be proved in parallel with the proof of Theorem 1, so the details of proofs are omitted. □

Corollary 2.2. *If we set,* $\alpha = \lambda = 1$ *and* $\mu = 3/2$ *in* (2.3) *then we obtain the solution of fractional kinetic equation involving Struve function* $H_{v}(z)$ *as : For* $d > 0, \upsilon > 0, c, l, t \in \mathbb{C}$, $\Re(l) > -1$ *and* $\Re(u) > 0$, *then the solution of the equation ,*

$$N(t) - N_0 H_{l,3/2}^{1,1}(d^{v}t^{v}) := -d^{v}\ {}_0D_t^{-v}N(t),$$

is given by

$$N(t) = N_0 \left(\frac{d^{v}}{2}\right)^{l+1} t^{lv+v-1} \times \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(2kv + vl + \nu + 1)}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+l+\frac{3}{2}\right)} \left(\frac{d^{v}t^{v}}{2}\right)^{2k} E_{v,(2k+l+1)v}(-d^{v}t^{v})$$

Theorem 3. *If* $d > 0, \upsilon > 0, a, c, b, l, t \in \mathbb{C}$, $\Re(l) > -1, \Re(u) > 0$ *with* $|u| < d^{-1}$ *and* $\mathfrak{a} \neq d$ *then for the solution of the equation*

$$N(t) - N_0 H_{l,\mu}^{\lambda,\alpha}(d^{v}t^{v}) = -\mathfrak{a}^{v}\,{}_0D_t^{-v}N(t), \tag{2.5}$$

there hold the formula

$$\begin{aligned} N(t) &= N_0 \left(\frac{d^{v}}{2}\right)^{l+1} t^{lv+v-1} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(2kv + vl + v + 1)}{\Gamma(\alpha k + \mu)\Gamma\left(\lambda k + l + \frac{3}{2}\right)} \\ &\quad \times \left(\frac{d^{v}t^{v}}{2}\right)^{2k} E_{v,(2k+l+1)v}(-\mathfrak{a}^{v}t^{v}), \end{aligned} \tag{2.6}$$

where $E_{v,(2k+l+1)v}(-\mathfrak{a}^{v}t^{v})$ *is the Mittag-Leffler function in* (1.6).

Proof. The proof of theorem 3 is derived similarly as that of theorems 1 and 2. □

Corollary 2.3. *By setting* $\alpha = \lambda = 1$ *and* $\mu = 3/2$ *in* (2.5) *we obtain the solution of fractional kinetic equations as : For* $d > 0, \upsilon > 0, c, l, t \in \mathbb{C}$,$\mathfrak{a} \neq d$, $\Re(l) > -1$ $\Re(u) > 0$ *and then the solution of the equation ,*

$$N(t) - N_0 H_{l,3/2}^{1,1}(d^{v}t^{v}) := -\mathfrak{a}^{v}\ {}_0D_t^{-v}N(t),$$

is given by

$$\begin{aligned} N(t) &= N_0 \left(\frac{d^{v}}{2}\right)^{l+1} t^{lv+v-1} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(2kv + vl + v + 1)}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+l+\frac{3}{2}\right)} \\ &\quad \times \left(\frac{d^{v}t^{v}}{2}\right)^{2k} E_{v,(2k+l+1)v}(-\mathfrak{a}^{v}t^{v}) \end{aligned}$$

2.1. **Special Cases.**

1 . Consider the generalized Struve function given by Bhowmic [11]

$$H_l^{\lambda}(x)=\sum_{k-0}^{\infty}\frac{(-1)^k\left(\frac{t}{2}\right)^{2k+l+1}}{\Gamma\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)} \tag{2.7}$$

Now, we have the following corollaries due to theorems 1, 2 and 3 respectively.

Corollary 2.4. *If $d>0,\upsilon>0,\lambda,l,t\in\mathbb{C}$, $\Re(l)>-1$ and $\Re(u)>0$ then the solution of the equation*

$$N(t)-N_0H_l^{\lambda}(t):=-d^{\upsilon}\ {}_0D_t^{-\upsilon}N(t),$$

is given by the following formula

$$N(t)=\frac{N_0}{2}\sum_{k-0}^{\infty}\frac{(-1)^k\Gamma(2k+l+2)}{\Gamma\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)}\left(\frac{t}{2}\right)^{2k+l}E_{\upsilon,2k+l+1}(-d^{\upsilon}t^{\upsilon})$$

Corollary 2.5. *If $d>0,\upsilon>0,\lambda,l,t\in\mathbb{C}$, $\Re(l)>-1$ and $\Re(u)>0$ then for the solution of the equation*

$$N(t)-N_0H_l^{\lambda}(d^{\upsilon}t^{\upsilon})=-d^{\upsilon}{}_0D_t^{-\upsilon}N(t)$$

there holds the formula

$$N(t)=N_0\left(\frac{d^{\upsilon}}{2}\right)^{l+1}t^{l\upsilon+\upsilon-1}\sum_{k=0}^{\infty}\frac{(-1)^k\Gamma(2k\upsilon+\upsilon l+\nu+1)}{\Gamma\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)}\left(\frac{d^{\upsilon}t^{\upsilon}}{2}\right)^{2k}E_{\upsilon,(2k+l+1)\upsilon}(-d^{\upsilon}t^{\upsilon})$$

Corollary 2.6. *If $d>0,\upsilon>0,\lambda,b,l,t\in\mathbb{C}$,$\mathfrak{a}\neq d$,$\Re(l)>-1$ and $\Re(u)>0$ then for the solution of the equation*

$$N(t)-N_0H_l^{\lambda}(d^{\upsilon}t^{\upsilon})=-\mathfrak{a}^{\upsilon}{}_0D_t^{-\upsilon}N(t)$$

there hold the formula

$$\begin{aligned}N(t) &= N_0\left(\frac{d^{\upsilon}}{2}\right)^{l+1}t^{l\upsilon+\upsilon-1}\sum_{k=0}^{\infty}\frac{(-1)^k\Gamma(2k\upsilon+\upsilon l+\upsilon+1)}{\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)}\\ &\times\left(\frac{d^{\upsilon}t^{\upsilon}}{2}\right)^{2k}E_{\upsilon,(2k+l+1)\upsilon}(-\mathfrak{a}^{\upsilon}t^{\upsilon})\end{aligned}$$

2 . Consider the generalized Struve function given by Kant [23]

$$H_l^{\lambda,\alpha}(x)=\sum_{k-0}^{\infty}\frac{(-1)^k\left(\frac{x}{2}\right)^{2k+l+1}}{\Gamma\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(\alpha k+\frac{3}{2}\right)} \tag{2.8}$$

Now, we have the following corollaries due to theorem 1, 2 and 3 respectively.

Corollary 2.7. *If* $d>0,\upsilon>0,\lambda,l,t\in\mathbb{C}$, $\Re(l)>-1$ *and* $\Re(u)>0$, *then the solution of the equation*

$$N(t)-N_0H_l^{\lambda,\alpha}(t):=-d^{\upsilon}\ {}_0D_t^{-\upsilon}N(t),$$

is given by the following formula

$$N(t)=\frac{N_0}{2}\sum_{k-0}^{\infty}\frac{(-1)^k\Gamma(2k+l+2)}{\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(\alpha k+\frac{3}{2}\right)}\left(\frac{t}{2}\right)^{2k+l}E_{\upsilon,2k+l+1}(-d^{\upsilon}t^{\upsilon})$$

Corollary 2.8. *If* $d>0,\upsilon>0,\lambda,l,t\in\mathbb{C}$,$\Re(l)>-1$ *and* $\Re(u)>0$ *then for the solution of the equation*

$$N(t)-N_0H_l^{\lambda,\alpha}(d^{\upsilon}t^{\upsilon})=-d^{\upsilon}{}_0D_t^{-\upsilon}N(t)$$

there holds the formula

$$N(t)=N_0\left(\frac{d^{\upsilon}}{2}\right)^{l+1}t^{l\upsilon+\upsilon-1}\sum_{k=0}^{\infty}\frac{(-1)^k\Gamma(2k\upsilon+\upsilon l+\nu+1)}{\Gamma\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(\alpha k+\frac{3}{2}\right)}\left(\frac{d^{\upsilon}t^{\upsilon}}{2}\right)^{2k}E_{\upsilon,(2k+l+1)\upsilon}(-d^{\upsilon}t^{\upsilon})$$

Corollary 2.9. *If* $d>0,\upsilon>0,\lambda,b,l,t\in\mathbb{C}$,$\mathfrak{a}\neq d$,$\Re(l)>-1$ *and* $\Re(u)>0$ *then for the solution of the equation*

$$N(t)-N_0H_l^{\lambda,\alpha}(d^{\upsilon}t^{\upsilon})=-\mathfrak{a}^{\upsilon}{}_0D_t^{-\upsilon}N(t)$$

there hold the formula

$$\begin{aligned}N(t) &= N_0\left(\frac{d^{\upsilon}}{2}\right)^{l+1}t^{l\upsilon+\upsilon-1}\sum_{k=0}^{\infty}\frac{(-1)^k\Gamma(2k\upsilon+\upsilon l+\upsilon+1)}{\left(\lambda k+l+\frac{3}{2}\right)\Gamma\left(\alpha k+\frac{3}{2}\right)}\\ &\times\left(\frac{d^{\upsilon}t^{\upsilon}}{2}\right)^{2k}E_{\upsilon,(2k+l+1)\upsilon}(-\mathfrak{a}^{\upsilon}t^{\upsilon})\end{aligned}$$

3 . Consider the generalized Struve function given by Singh [37]

$$H_{l,\mu}^{\lambda}(x)=\sum_{k-0}^{\infty}\frac{(-1)^k\left(\frac{x}{2}\right)^{2k+l+1}}{\Gamma\left(\lambda k+\frac{l}{\mu}+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)}\tag{2.9}$$

Now, we have the following corollaries due to theorem 1, theorem 2 and theorem 3 respectively.

Corollary 2.10. *If* $d>0,\upsilon>0,\lambda,l,t\in\mathbb{C}$, $\Re(l)>-1$ *and* $\Re(u)>0$ *then the solution of the equation*

$$N(t)-N_0H_{l,\mu}^{\lambda}(t):=-d^{\upsilon}\ {}_0D_t^{-\upsilon}N(t),$$

is given by the following formula

$$N(t)=\frac{N_0}{2}\sum_{k-0}^{\infty}\frac{(-1)^k\Gamma(2k+l+2)}{\Gamma\left(\lambda k+\frac{l}{\mu}+\frac{3}{2}\right)\Gamma\left(k+\frac{3}{2}\right)}\left(\frac{t}{2}\right)^{2k+l}E_{\upsilon,2k+l+1}(-d^{\upsilon}t^{\upsilon})$$

Corollary 2.11. *If* $d > 0, \upsilon > 0, \lambda, l, t \in \mathbb{C}$, $\Re(l) > -1$ *and* $\Re(u) > 0$ *then for the solution of the equation*

$$N(t) - N_0 H_{l,\mu}^{\lambda}(d^{\upsilon} t^{\upsilon}) = -d^{\upsilon}\, {}_0D_t^{-\upsilon} N(t)$$

there holds the formula

$$N(t) = N_0 \left(\frac{d^{\upsilon}}{2}\right)^{l+1} t^{l\upsilon+\upsilon-1} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(2k\upsilon + \upsilon l + \nu + 1)}{\Gamma\left(\lambda k + \frac{l}{\mu} + \frac{3}{2}\right) \Gamma\left(k + \frac{3}{2}\right)} \left(\frac{d^{\upsilon} t^{\upsilon}}{2}\right)^{2k} E_{\upsilon,(2k+l+1)\upsilon}(-d^{\upsilon} t^{\upsilon}).$$

Corollary 2.12. *If* $d > 0, \upsilon > 0, \lambda, b, l, t \in \mathbb{C}$, $\mathfrak{a} \neq d$, $\Re(l) > -1, \mu$ *is an arbitrary parameter and* $\Re(u) > 0$ *then for the solution of the equation*

$$N(t) - N_0 H_{l,\mu}^{\lambda}(d^{\upsilon} t^{\upsilon}) = -\mathfrak{a}^{\upsilon}\, {}_0D_t^{-\upsilon} N(t)$$

there hold the formula

$$\begin{aligned} N(t) &= N_0 \left(\frac{d^{\upsilon}}{2}\right)^{l+1} t^{l\upsilon+\upsilon-1} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(2k\upsilon + \upsilon l + \upsilon + 1)}{\left(\lambda k + \frac{l}{\mu} + \frac{3}{2}\right) \Gamma\left(k + \frac{3}{2}\right)} \\ &\times \left(\frac{d^{\upsilon} t^{\upsilon}}{2}\right)^{2k} E_{\upsilon,(2k+l+1)\upsilon}(-\mathfrak{a}^{\upsilon} t^{\upsilon}) \end{aligned}$$

References

[1] A. R. Ahmadi, S.E. Widnall, *Unsteady lifting-line theory as a singular-perturbation problem*, J. Fluid Mech. **153** (1985) 59–81.

[2] M.A. Asiru, *Sumudu transform and the solution of integral equation of convolution type*, International Journal of Mathematical Education in Science and Technology, **32** (2001) 906–910.

[3] F.B.M. Belgacem , A.A. Karaballi, S.L., Kalla, *Analytical Investigations of the Sumudu Transform and Applications to Integral Production Equations*, Journal of Mathematical Problems in Engineering, **3** 2003, 103–118.

[4] F.B.M. Belgacem, A.A. Karaballi, *Sumudu transform fundamental properties investigations and applications*, International J. Appl. Math. Stoch. Anal. (2005) 1–23.

[5] F.B.M. Belgacem , A.A.Karaballi, *Sumudu transform fundamental properties investigations and applications*, Journal of Applied Mathematics and Stochastic Analysis, **2006** (2006), Article ID 91083.

[6] F.B.M. Belgacem, *Introducing and analyzing deeper Sumudu properties*, Nonlinear Studies, **13** (2006) 23–42.

[7] F.B.M. Belgacem, *Applications with the Sumudu transform to Bessel functions and equations*, Applied Math. Sciences, **4** (2010) 3665–3686.

[8] F.B.M. Belgacem, *Sumudu applications to Maxwells equations*, PIERS Online, **5** (2009) 355–360.

[9] F.B.M. Belgacem, E.H., Al-Shemas E. H., R. Silambarasan, *Sumudu computation of the transient magnetic field in a lossy medium.* Applied Mathematics & Information Sciences, **6** (2016) 1-9.

[10] F.B.M. Belgacem, R. Silambarasan, *A distinctive Sumudu treatment of trigonometric functions*, Journal of Computational and Applied Mathematics, **312** (2017) 74-81.

[11] K. N. Bhowmick, *Some relations between a generalized Struve's function and hypergeometric functions*, Vijnana Parishad Anusandhan Patrika, **5** (1962) 93–99.

[12] K. N. Bhowmick, *A generalized Struve's function and its recurrence formula*, Vijnana Parishad Anusandhan Patrika, **6** (1963) 1–11.

[13] V.B.L., Chaurasia, R.S., Dubey, F.B.M., Belgacem, *Fractional radial diffusion equation analytical solution via Hankel and Sumudu transforms*, International Journal of Mathematics in Eng. Science, & Aerospace, **3**2 (2012) 179–188.

[14] V.B.L. Chaurasia, S.C. Pandey, *On the new computable solution of the generalized fractional kinetic equations involving the generalized function for the fractional calculus and related functions*, Astrophys. Space Sci.**317** (2008) 213–219.

[15] A. Chouhan, S. Sarswat, *On solution of generalized Kinetic equation of fractional order*, Int. Jr. of Mathematical Sciences and Applications **2**2 (2012) 813–818.

[16] A. Chouhan, S.D. Purohit, S. Saraswat, *An alternative method for solving generalized differential equations of fractional order*, Kragujevac J. Math. **37**2 (2013) 299–306.

[17] A. Erdélyi, W. Magnus, F. Oberhettinger, F.G. Tricomi, *Tables of Integral Transforms*, Vol. 1, McGraw-Hill, New York-Toronto-London, (1954).

[18] V.G. Gupta, B. Sharma, F.B.M, Belgacem, *On the solutions of generalized fractional kinetic equations*, Applied Mathematical Sciences **5**19 (2011) 899–910.

[19] A. Gupta, C.L. Parihar, *On solutions of generalized kinetic equations of fractional order*, Bol. Soc. Paran. Mat. **32**1 (2014) 181–189.

[20] H.J. Haubold, A.M. Mathai, *The fractional kinetic equation and thermonuclear functions*, Astrophys. Space Sci. **327** (2000) 53–63.

[21] M .H. Hirata, *Flow near the bow of a steadily turning ship*, J. Fluid Mech. **71**2 (1975) 283–291.

[22] M.G.M., Hussain, F.B.M.,Belgacem, *Transient Solutions of Maxwells Equations Based on Sumudu Transformation*, Journal of Progress in Electromagnetic Research (PIER) **74** (2007) 273–289.

[23] B. N. Kanth, *Integrals involving generalized Struve's function*, Nepali Math. Sci. Rep. **6** (1981) 61–64.

[24] G.M. Mittag-Leffler, *Mittag-Leffler, Sur la representation d'une fonction monogene cinquieme note*, Acta Math.**29** (1905),101–181.

[25] K. S. Nisar, D. Baleanu, and M. M. A. Qurashi, *Fractional calculus and application of generalized Struve function*, SpringerPlus **5** (2016) 910 .

[26] K. S. Nisar, J. Choi, *Generalized Fractional Kinetic Equations Involving the generalized modified k-Bessel function*, arXiv:1612.03267 [math.CA], (2016)

[27] K.S. Nisar, S.D. Purohit, Saiful. R. Mondal, *Generalized fractional kinetic equations involving generalized Struve function of the first kind*, Journal of King Saud University Science **28** 2 (2016) 167–171

[28] K.S. Nisar, S.R. Mondal, F.B.M. Belgacem, *On Solution of fractional kinetic equation Involving k-Struve function*, arXiv:1611.09154 [math.CA], (2016)

[29] T. G. Pedersen, *Variational approach to excitons in carbon nanotubes*, Phys. Rev. B **67** (2003) 0734011–0734014.

[30] A. Saichev, M. Zaslavsky, *Fractional kinetic equations: solutions and applications*, Chaos, **7** (1997) 753-764.

[31] R.K. Saxena, A.M. Mathai, H.J. Haubold, *On fractional kinetic equations*, Astrophys. Space Sci. **282** (2002), 281-287.

[32] R.K. Saxena, A.M. Mathai, H.J. Haubold, *On generalized fractional kinetic equations*, Physica A 344 (2004) 657–664.

[33] R.K. Saxena, A.M. Mathai, H.J. Haubold, *Solution of generalized fractional reaction-diffusion equations*, Astrophys. Space Sci. **305** (2006) 305–313.

[34] R.K. Saxena and S.L. Kalla, *On the solutions of certain fractional kinetic equations*, Appl. Math. Comput. **199** (2008) 504–511.

[35] J. Shao, P. Hanggi, *Decoherent dynamics of a two-level system coupled to a sea of spins*, Phys. Rev. Lett.81 , 26, 5710-5713,(1998).

[36] D. C. Shaw, *Perturbational results for di raction of water-waves by nearly-vertical barriers*, IMA, J. Appl. Math. **34** (1985) 99–117.

[37] R. P. Singh, *Generalized Struve's function and its recurrence equation*, Vijnana Parishad Anusandhan Patrika **28**3 (1985) 287–292.

[38] M.R. Spiegel, *Theory and problems of Laplace transforms*, Schaums Outline Series, McGraw-Hill, New York, (1965).

[39] H.M. Srivastava, R.K. Saxena, *Operators of fractional integration and their applications*, Appl. Math. Comput.**118** (2001) 1–52.

[40] H. Struve, *Beitrag zur Theorie der Diffraction an Fernröhren*, Ann. Physik Chemie **17** (1882) 1008–1016.

[41] G.K. Watugala, *Sumudu transform:a new integral transform to solve differential equations and control engineering problems*, Int.J.Math. Edu.Sci.Tech. **24** (1993) 35–43.

[42] G. K.Watugala, *The Sumudu transform for functions of two variables*, Math. Eng.Ind. **8** (2002) 293–302.

[43] G.M. Zaslavsky, *Fractional kinetic equation for Hamiltonian chaos*, Physica D, **76** (1994) 110–122.

K. S. Nisar : Department of Mathematics, College of Arts and Science, Prince Sattam bin Abdulaziz University, Wadi Aldawaser, Riyadh region 11991, Saudi Arabia

E-mail address: ksnisar1@gmail.com

F.B.M. Belgacem : Department of Mathematics, Faculty of Basic Education, PAAET, Al-Ardhiya, Kuwait

E-mail address: fbmbelgacem@gmail.com

M. S. Abouzaid : Department of Mathematics, Faculty of Science, Kafrelshiekh University, Egypt.

E-mail address: moheb_abouzaid@hotmail.com